

Large- x structure of physical evolution kernels in Deep Inelastic Scattering

G. Grunberg

*Centre de Physique Théorique, École Polytechnique
91128 Palaiseau Cedex, France
E-mail: georges.grunberg@cpht.polytechnique.fr*

ABSTRACT: The modified evolution equation for parton distributions of Dokshitzer, Marchesini and Salam is extended to non-singlet Deep Inelastic Scattering coefficient functions and the physical evolution kernels which govern their scaling violation. Considering the $x \rightarrow 1$ limit, it is found that the leading next-to-eikonal logarithmic contributions to the physical kernels at any loop order can be expressed in term of the one-loop cusp anomalous dimension, a result which can presumably be extended to all orders in $(1 - x)$, and has eluded so far threshold resummation. Similar results are shown to hold for fragmentation functions in semi-inclusive e^+e^- annihilation. Gribov-Lipatov relation is found to be satisfied by the leading logarithmic part of the modified physical evolution kernels.

Contents

1. Introduction	1
2. The modified physical kernel	3
3. Leading next-to-eikonal logarithms	5
3.1 Two loop kernel	5
3.2 Three loop kernel	5
3.3 Four loop kernel	6
3.4 Five loop kernel	6
3.5 All-order relations	6
4. Leading next-to-next-to-eikonal logarithms	7
5. Fragmentation functions in e^+e^- annihilation	8
6. Conclusion	9

1. Introduction

There has been recently renewed interest [1–9] in threshold resummation of “next-to-eikonal” logarithmically enhanced terms which are suppressed by some power of the gluon energy $(1-x)$ for $x \rightarrow 1$ in momentum space (or by some power of $1/N$, $N \rightarrow \infty$ in moment space). In particular, in [2, 5–9] this question has been investigated at the level of “physical evolution kernels” which control the scaling violation of (non-singlet) structure functions. The scale-dependence of the Deep Inelastic Scattering (DIS) coefficient function $\mathcal{C}_2(x, Q^2, \mu_F^2)$ corresponding to the flavor non-singlet $F_2(x, Q^2)$ structure function ($F_2(x, Q^2)/x = \mathcal{C}_2(x, Q^2, \mu_F^2) \otimes q_{2,ns}(x, \mu_F^2)$, where $q_{2,ns}(x, \mu_F^2)$ is the corresponding quark distribution) can be expressed in terms of $\mathcal{C}_2(x, Q^2, \mu_F^2)$ itself, yielding the following “physical” evolution equation (see e.g. Refs. [10–14]):

$$\frac{\partial \mathcal{C}_2(x, Q^2, \mu_F^2)}{\partial \ln Q^2} = \int_x^1 \frac{dz}{z} K(z, a_s(Q^2)) \mathcal{C}_2(x/z, Q^2, \mu_F^2) \equiv K(x, a_s(Q^2)) \otimes \mathcal{C}_2(x, Q^2, \mu_F^2) , \quad (1.1)$$

where μ_F is the factorization scale (I assume for definiteness the $\overline{MS}$ factorization scheme is used). $K(x, a_s(Q^2))$ is the momentum space *physical evolution kernel*, or *physical anomalous dimension*; it is independent of the factorization scale and renormalization-scheme invariant.

In [15], the result for the leading contribution to this quantity in the $x \rightarrow 1$ limit was derived, which resums all logarithms at the leading eikonal level, and nicely summarizes analytically in *momentum* space the standard results [16, 17] of threshold resummation:

$$K(x, a_s(Q^2)) \sim \frac{\mathcal{J}(rQ^2)}{r} + B_\delta^{DIS}(a_s(Q^2)) \delta(1-x), \quad (1.2)$$

where $r = \frac{1-x}{x}$ (with $rQ^2 \equiv W^2$ the final state “jet” mass), $B_\delta^{DIS}(a_s)$ is related to the quark form factor, and $\mathcal{J}(Q^2)$, the “physical Sudakov anomalous dimension” (a renormalization scheme invariant quantity), is given by:

$$\begin{aligned} \mathcal{J}(Q^2) &= A(a_s(Q^2)) + \frac{dB(a_s(Q^2))}{d \ln Q^2} \\ &= A(a_s(Q^2)) + \beta(a_s(Q^2)) \frac{dB(a_s(Q^2))}{da_s} \equiv \sum_{i=1}^{\infty} j_i a_s^i(Q^2). \end{aligned} \quad (1.3)$$

In eq.(1.3),

$$A(a_s) = \sum_{i=1}^{\infty} A_i a_s^i \quad (1.4)$$

is the universal “cusp” anomalous dimension [18] (see also [19]), with $a_s \equiv \frac{\alpha_s}{4\pi}$ the $\overline{MS}$ coupling,

$$\beta(a_s) = \frac{da_s}{d \ln Q^2} = -\beta_0 a_s^2 - \beta_1 a_s^3 - \beta_2 a_s^4 + \dots \quad (1.5)$$

is the beta function (with $\beta_0 = \frac{11}{3}C_A - \frac{2}{3}n_f$) and

$$B(a_s) = \sum_{i=1}^{\infty} B_i a_s^i \quad (1.6)$$

is the usual final state “jet function” anomalous dimension. It should be noted that $j_1 = A_1$ (the one loop cusp anomalous dimension), and also that both $A(a_s)$ and $B(a_s)$ (in contrast to $\mathcal{J}(Q^2)$) are renormalization scheme-dependent quantities. The renormalization group invariance of $\mathcal{J}(Q^2)$ yields the standard relation:

$$\begin{aligned} \mathcal{J}((1-x)Q^2) &= j_1 a_s + a_s^2 [-j_1 \beta_0 L_x + j_2] \\ &+ a_s^3 [j_1 \beta_0^2 L_x^2 - (j_1 \beta_1 + 2j_2 \beta_0) L_x + j_3] \\ &+ a_s^4 [-j_1 \beta_0^3 L_x^3 + (\frac{5}{2} j_1 \beta_1 \beta_0 + 3j_2 \beta_0^2) L_x^2 - (j_1 \beta_2 + 2j_2 \beta_1 + 3j_3 \beta_0) L_x + j_4] + \dots, \end{aligned} \quad (1.7)$$

where $L_x \equiv \ln(1-x)$ and $a_s = a_s(Q^2)$, from which the structure of all the eikonal logarithms in $K(x, a_s(Q^2))$ can be derived. A term like $\frac{L_x^p}{1-x}$ arising from $\frac{\mathcal{J}(rQ^2)}{r}$ in eq.(1.2) must be interpreted as usual as a standard $+$ -distribution. *All* the eikonal logarithms are thus absorbed into the *single* scale $(1-x)Q^2$ (see also [20–22] and section VI-E in [11]).

However, no analogous result holds [6] at the next-to-eikonal level (except [2] at large- β_0). In this note, I show that the *leading* next-to-eikonal logarithmic contributions to the physical evolution kernel at a given order in a_s can actually be determined in term of lower order *leading* eikonal coefficients, representing the first step towards threshold resummation at the next-to-eikonal level. This result is obtained by extending the approach of [23, 24] (which deals with parton distributions) to the DIS coefficient functions themselves.

2. The modified physical kernel

I consider the class of modified physical evolution equations:

$$\frac{\partial \mathcal{C}_2(x, Q^2, \mu_F^2)}{\partial \ln Q^2} = \int_x^1 \frac{dz}{z} K(z, a_s(Q^2), \lambda) \mathcal{C}_2(x/z, Q^2/z^\lambda, \mu_F^2) , \quad (2.1)$$

where for book-keeping purposes I introduced the parameter λ , which shall eventually be set to its physically meaningful value $\lambda = 1$, in straightforward analogy to the modified evolution equation for parton distributions of [24]. I note that $K(x, a_s, \lambda = 0) \equiv K(x, a_s)$, the ‘standard’ physical evolution kernel. Eq.(2.1) allows to determine $K(x, a_s, \lambda)$ given $K(x, a_s)$ (or vice-versa). Indeed, expanding $\mathcal{C}_2(y, Q^2/z^\lambda, \mu_F^2)$ around $z = 1$, keeping the other two variables fixed, and reporting into eq.(2.1), one easily derives the following relation between $K(x, a_s, \lambda)$ and $K(x, a_s)$:

$$\begin{aligned} K(x, a_s) &= K(x, a_s, \lambda) - \lambda [\ln x \, K(x, a_s, \lambda)] \otimes K(x, a_s) \\ &\quad + \frac{\lambda^2}{2} [\ln^2 x \, K(x, a_s, \lambda)] \otimes [\beta(a_s) \frac{\partial K(x, a_s)}{\partial a_s} + K(x, a_s) \otimes K(x, a_s)] + \dots , \end{aligned} \quad (2.2)$$

where only terms with a single overall factor of λ need actually to be kept up to next-to-eikonal order, since one can check terms with more factors of λ , which are associated to more factors of $\ln x$, are not relevant to determine the next-to-eikonal logarithms in the physical kernel. In the rest of the paper (except section 4) I shall therefore simply use:

$$K(x, a_s) = K(x, a_s, \lambda) - \lambda [\ln x \, K(x, a_s, \lambda)] \otimes K(x, a_s) + \dots . \quad (2.3)$$

Eq.(2.3) can be solved perturbatively. Setting:

$$K(x, a_s, \lambda) = K_0(x, \lambda) a_s + K_1(x, \lambda) a_s^2 + K_2(x, \lambda) a_s^3 + K_3(x, \lambda) a_s^4 + \dots \quad (2.4)$$

(and similarly for $K(x, a_s)$), one gets:

$$\begin{aligned} K_0(x, \lambda) &= K_0(x) \\ K_1(x, \lambda) &= K_1(x) + \lambda [\ln x \, K_0(x)] \otimes K_0(x) \\ K_2(x, \lambda) &= K_2(x) + \lambda \{ [\ln x \, K_1(x)] \otimes K_0(x) + [\ln x \, K_0(x)] \otimes K_1(x) \} + \dots \\ K_3(x, \lambda) &= K_3(x) + \lambda \{ [\ln x \, K_2(x)] \otimes K_0(x) + [\ln x \, K_1(x)] \otimes K_1(x) \\ &\quad + [\ln x \, K_0(x)] \otimes K_2(x) \} + \dots . \end{aligned} \quad (2.5)$$

The $K_i(x)$'s are determined in term of splitting functions and coefficient functions as follows [14]:

$$\begin{aligned} K_0(x) &= P_0(x) \\ K_1(x) &= P_1(x) - \beta_0 c_1(x) \\ K_2(x) &= P_2(x) - \beta_1 c_1(x) - \beta_0(2c_2(x) - c_1^{\otimes 2}(x)) \\ K_3(x) &= P_3(x) - \beta_2 c_1(x) - \beta_1(2c_2(x) - c_1^{\otimes 2}(x)) - \beta_0(3c_3(x) - 3c_2(x) \otimes c_1(x) + c_1^{\otimes 3}(x)) , \end{aligned} \quad (2.6)$$

where $P_i(x)$ are the standard $(i+1)$ -loop splitting functions, $c_i(x)$ are the i -loop coefficient functions, and $c_1^{\otimes 2}(x) \equiv c_1(x) \otimes c_1(x)$, etc....

Consider now the $x \rightarrow 1$ limit. The one-loop splitting function is given by [25]:

$$P_0(x) = A_1 p_{qq}(x) + B_1^\delta \delta(1-x) , \quad (2.7)$$

with $A_1 = 4C_F$, and¹:

$$p_{qq}(x) = \frac{1}{1-x} - 1 + \frac{1}{2}(1-x) = \frac{x}{1-x} + \frac{1}{2}(1-x) = \frac{1}{r} + \frac{1}{2}(1-x) . \quad (2.8)$$

Moreover, at the next-to-eikonal level we have, dropping from now on δ function contributions:

$$P_1(x) = \frac{A_2}{r} + C_2 L_x + D_2 + \dots , \quad (2.9)$$

with [26]:

$$C_2 = A_1^2 . \quad (2.10)$$

Also:

$$c_1(x) = \frac{c_{11} L_x + c_{10}}{r} + b_{11} L_x + b_{10} + \dots \quad (2.11)$$

with $c_{11} = A_1 = 4C_F$, $b_{11} = 0$. From eq.(2.6) one can derive [6,7] the following expansions for $x \rightarrow 1$:

$$\begin{aligned} K_0(x) &= P_0(x) = \frac{k_{10}}{r} + h_{10} + \dots \\ K_1(x) &= \frac{k_{21} L_x + k_{20}}{r} + h_{21} L_x + h_{20} + \dots \\ K_2(x) &= \frac{k_{32} L_x^2 + k_{31} L_x + k_{30}}{r} + h_{32} L_x^2 + h_{31} L_x + h_{30} + \dots \\ K_3(x) &= \frac{k_{43} L_x^3 + k_{42} L_x^2 + k_{41} L_x + k_{40}}{r} + h_{43} L_x^3 + h_{42} L_x^2 + h_{41} L_x + h_{40} + \dots . \end{aligned} \quad (2.12)$$

¹ $p_{qq}(x)$ is defined to be 1/2 the corresponding function in [7].

3. Leading next-to-eikonal logarithms

3.1 Two loop kernel

From eq.(2.6), (2.7), (2.9) and (2.11) one deduces: $k_{10} = A_1$, $h_{10} = 0$, and $k_{21} = -\beta_0 A_1$, $h_{21} = C_2$. Then eq.(2.5) yields for $x \rightarrow 1$:

$$\begin{aligned} K_0(x, \lambda) &= P_0(x) \\ K_1(x, \lambda) &= \frac{k_{21}L_x + k_{20}}{r} + (h_{21} - \lambda k_{10}^2)L_x + \mathcal{O}(L_x^0) . \end{aligned} \quad (3.1)$$

Now

$$h_{21}(\lambda) = h_{21} - \lambda k_{10}^2 = C_2 - \lambda A_1^2 = (1 - \lambda)A_1^2 . \quad (3.2)$$

Thus, setting $\lambda = 1$, one finds that the leading next-to-eikonal logarithm in $K_1(x, \lambda = 1)$ vanishes, yielding the relation:

$$h_{21} = k_{10}^2 = 16C_F^2 , \quad (3.3)$$

which is correct [6, 7]. This finding is not surprising: up to two loop, the leading next-to-eikonal logarithm is contributed only by the splitting function, since $b_{11} = 0$ (e.g. $h_{21} = C_2$), and one effectively recovers the result (eq.(2.10)) holding [24] for the two loop splitting function. The situation however changes drastically at three loop, where the leading next-to-eikonal logarithm is contributed by the coefficient function rather than the splitting function, and the crucial question is whether the leading next-to-eikonal logarithm still vanishes for $\lambda = 1$.

3.2 Three loop kernel

Eq.(2.5) yields for $x \rightarrow 1$:

$$K_2(x, \lambda) = \frac{k_{32}L_x^2 + k_{31}L_x + k_{30}}{r} + (h_{32} - \lambda \frac{3}{2}k_{21}k_{10})L_x^2 + \mathcal{O}(L_x) . \quad (3.4)$$

Requiring $h_{32}(\lambda)$, the coefficient of the $\mathcal{O}(L_x^2)$ term, to vanish for $\lambda = 1$ predicts:

$$h_{32} = \frac{3}{2}k_{21}k_{10} = -\frac{3}{2}\beta_0 A_1^2 = -24\beta_0 C_F^2 , \quad (3.5)$$

which is indeed the correct [6, 7] value. I stress that this result is *not* a consequence of the relation [24, 27, 28] $C_3 = 2A_1A_2$ for $P_2(x)$. Indeed it is well-known [29] that the $P_i(x)$'s, and in particular $P_2(x)$, have only a *single* next-to-eikonal logarithm:

$$P_2(x) = \frac{A_3}{r} + C_3L_x + D_3 + \dots , \quad (3.6)$$

and thus $P_2(x)$ cannot contribute to the *double* logarithm in $K_2(x)$. Rather, h_{32} is contributed by the coefficient functions in eq.(2.6), and eq.(3.5) yields a prediction for the $\mathcal{O}(L_x^2)$ term in $c_2(x)$.

3.3 Four loop kernel

Eq.(2.5) yields for $x \rightarrow 1$:

$$K_3(x, \lambda) = \frac{k_{43}L_x^3 + k_{42}L_x^2 + k_{41}L_x + k_{40}}{r} + [h_{43} - \lambda(\frac{4}{3}k_{10}k_{32} + \frac{1}{2}k_{21}^2)]L_x^3 + \mathcal{O}(L_x^2) , \quad (3.7)$$

where $k_{32} = A_1\beta_0^2$ (consistently with eq.(1.7)). Requiring $h_{43}(\lambda)$, the coefficient of the $\mathcal{O}(L_x^3)$ term, to vanish for $\lambda = 1$ predicts:

$$h_{43} = \frac{4}{3}k_{10}k_{32} + \frac{1}{2}k_{21}^2 = \frac{11}{6}\beta_0^2A_1^2 = \frac{88}{3}\beta_0^2C_F^2 , \quad (3.8)$$

which is again the correct [6, 7] value.

3.4 Five loop kernel

One can similarly predict the leading next-to-eikonal logarithm in the five loop physical kernel (which depends on the four loop coefficient function). Using eq.(2.3), the coefficient of the $\mathcal{O}(L_x^4)$ term in $K_4(x, \lambda)$ is found to be given by:

$$h_{54}(\lambda) = h_{54} - \lambda(\frac{5}{4}k_{10}k_{43} + \frac{5}{6}k_{21}k_{32}) , \quad (3.9)$$

where $k_{43} = -A_1\beta_0^3$ (again consistent with eq.(1.7)). Requiring this coefficient to vanish for $\lambda = 1$ predicts²:

$$h_{54} = \frac{5}{4}k_{10}k_{43} + \frac{5}{6}k_{21}k_{32} = -\frac{25}{12}\beta_0^3A_1^2 = -\frac{100}{3}\beta_0^3C_F^2 . \quad (3.10)$$

3.5 All-order relations

Defining moments by

$$K(N, a_s) = \int_0^1 dx x^{N-1} K(x, a_s) , \quad (3.11)$$

eq.(2.3) yields in moment space:

$$K(N, a_s) = \frac{K(N, a_s, \lambda)}{1 + \lambda \dot{K}(N, a_s, \lambda)} , \quad (3.12)$$

where $\dot{f} \equiv \partial f / \partial N$. Assuming the leading next-to-eikonal logarithms vanish to all orders (in an expansion in $1/r$) for $\lambda = 1$, i.e. that $h_{i+1,i}(\lambda = 1) = 0$ for $i \geq 0$, one can derive [30] from eq.(3.12) the resummation formula:

$$\sum_{i=0}^{\infty} h_{i+1,i} L_x^i a_s^{i+1} = \frac{A_1}{\beta_0} \frac{A_1 a_s}{1 + a_s \beta_0 L_x} \ln(1 + a_s \beta_0 L_x) , \quad (3.13)$$

which correctly reproduces the results in the previous subsections.

²Hence $\xi_{\text{DIS}_4} = \frac{100}{3}$ in the notation of [7].

One can further show [30] that the moment space functional relation which accounts for *leading* logarithms at all orders in $(1-x)$ is:

$$K(N, a_s) = K[N - \lambda K(N, a_s), a_s, \lambda] . \quad (3.14)$$

Eq.(3.12) results from expanding the right hand side of eq.(3.14) to first order in $\Delta N \equiv \lambda K(N, a_s)$. It is interesting that eq.(3.14) is identical to the functional relation³ obtained [28, 32] for the splitting functions in the conformal limit (where the splitting functions coincide with the K_i 's).

4. Leading next-to-next-to-eikonal logarithms

It can be checked [30] that similar methods allow to predict using eq.(2.2) the *leading* logarithmic contributions at the next-to-next-to-eikonal level, i.e. the coefficient of the $(1-x)L_x^i$ term in $K_i(x)$. The crucial new point, however, is that the leading term in the eikonal expansion has to be defined in term of the one-loop splitting function prefactor $p_{qq}(x)$ (eq.(2.8)), instead of $1/r$ as in eq.(2.12). Namely, keeping only leading logarithms at each eikonal order, the predicted f_{ji}^c coefficients ($j = i + 1, i \geq 0$) are defined⁴ by:

$$K_i(x)|_{\text{LL}} = L_x^i [p_{qq}(x) k_{ji} + h_{ji} + (1-x)f_{ji}^c + (1-x)^2 g_{ji} + \mathcal{O}((1-x)^3)] . \quad (4.1)$$

Eq.(2.2) yields the corresponding $f_{ji}^c(\lambda)$ coefficients in $K_i(x, \lambda)$:

$$\begin{aligned} f_{21}^c(\lambda) &= f_{21}^c + \lambda \frac{1}{2} k_{10}^2 \\ f_{32}^c(\lambda) &= f_{32}^c - \lambda \left(-\frac{3}{4} k_{10} k_{21} + k_{10} h_{21} \right) + \lambda^2 \frac{1}{2} k_{10}^3 \\ f_{43}^c(\lambda) &= f_{43}^c - \lambda \left(-\frac{2}{3} k_{10} k_{32} + \frac{1}{2} (h_{21} - \frac{1}{2} k_{21}) k_{21} + k_{10} h_{32} \right) + \lambda^2 k_{10}^2 k_{21} , \end{aligned} \quad (4.2)$$

where I used that $h_{10} = 0$, and one should note the presence of contributions *quadratic* in λ . Assuming the $f_{ji}^c(\lambda)$'s vanish for $\lambda = 1$, one thus derives the relations (with $f_{10}^c = 0$):

$$\begin{aligned} f_{21}^c &= -\frac{1}{2} k_{10}^2 = -8C_F^2 \\ f_{32}^c &= -\frac{3}{4} k_{10} k_{21} + k_{10} h_{21} - \frac{1}{2} k_{10}^3 = 12C_F^2 \beta_0 + 32C_F^3 \\ f_{43}^c &= -\frac{2}{3} k_{10} k_{32} + \frac{1}{2} (h_{21} - \frac{1}{2} k_{21}) k_{21} + k_{10} h_{32} - k_{10}^2 k_{21} = -\frac{44}{3} C_F^2 \beta_0^2 - 64C_F^3 \beta_0 , \end{aligned} \quad (4.3)$$

which are seen to be correct using eq.(3.26) in [7]. The latter equation also makes it likely that similar leading logarithmic predictions can be obtained to any order in $(1-x)$, using

³A similar functional relation has been obtained in a different context in [31].

⁴The motivation for the superscript “c” (for “classical”) shall be clarified in the Conclusion section.

the *same* prefactor $p_{qq}(x)$ as in eq.(4.1) to define the leading term in the eikonal expansion. Indeed, one derives for instance [30] the $\mathcal{O}((1-x)^2)$ coefficients in eq.(4.1) (with $g_{10} = 0$):

$$\begin{aligned} g_{21} &= \frac{1}{3}k_{10}^2 = \frac{16}{3}C_F^2 \\ g_{32} &= \frac{1}{6}k_{10}k_{21} + (f_{21}^c + \frac{1}{2}h_{21} + \frac{1}{3}k_{21})k_{10} = \frac{1}{2}k_{10}k_{21} = -8C_F^2\beta_0, \end{aligned} \quad (4.4)$$

which are correct [7]. I note that f_{21}^c and g_{21} coincide (like h_{21}) with the splitting functions contributions.

5. Fragmentation functions in e^+e^- annihilation

Similar results hold for physical evolution kernels associated to fragmentation functions in semi-inclusive e^+e^- annihilation (SIA), provided one sets $\lambda = -1$ in the analogue of eq.(2.1):

$$\frac{\partial \mathcal{C}_T(x, Q^2, \mu_F^2)}{\partial \ln Q^2} = \int_x^1 \frac{dz}{z} K_T(z, a_s(Q^2), \lambda) \mathcal{C}_T(x/z, Q^2/z^\lambda, \mu_F^2), \quad (5.1)$$

where $\mathcal{C}_T$ denotes a generic non-singlet SIA coefficient function. I first note that threshold resummation in this case [33] leads at the leading eikonal level to an equation similar to eq.(1.2):

$$K_T(x, a_s(Q^2)) \sim \frac{\mathcal{J}((1-x)Q^2)}{1-x} + B_\delta^{SIA}(a_s(Q^2))\delta(1-x), \quad (5.2)$$

where x should now be identified to Feynman- x rather than Bjorken- x , and I used the results of [34] which imply that the “physical Sudakov anomalous dimension” $\mathcal{J}(Q^2)$ is *the same* for structure and fragmentation functions. The statement above eq.(5.1) then follows from the following two observations:

- i) The predictions in eq.(3.3), (3.5), (3.8) and (3.10) depend only upon coefficients of leading *eikonal* logarithms in the physical evolution kernels.
- ii) Eq.(3.26) in [7] shows that the latter coefficients are *identical* for deep-inelastic structure functions and for e^+e^- fragmentation functions (consistently with the remark below eq.(5.2)), but that the coefficients of the leading *next-to-eikonal* logarithms are equal only up to a sign change (in an expansion in $1/r$) between deep-inelastic structure functions and fragmentation functions.

One deduces the resummation formula ($j = i + 1$):

$$\sum_{i=0}^{\infty} h_{ji}^{SIA} L_x^i a_s^{i+1} = -\frac{A_1}{\beta_0} \frac{A_1 a_s}{1 + a_s \beta_0 L_x} \ln(1 + a_s \beta_0 L_x). \quad (5.3)$$

6. Conclusion

A modified⁵ evolution equation for DIS non-singlet structure functions, analogous to the one used in [24] for parton distributions, but which deals with the *physical* scaling violation and coefficient functions, has been proposed. It allows to relate the leading next-to-eikonal logarithmic contributions in the momentum space physical evolution kernel to coefficients of leading eikonal logarithms at lower loop order (depending only upon the one-loop cusp anomalous dimension A_1), which represents the first step towards threshold resummation at the next-to-eikonal level. This result also explains the observed [6, 7] *universality* of the leading next-to-eikonal logarithmic contributions to the physical kernels of the various non-singlet structure functions, linking them to the known [35] universality of the eikonal contributions. Similar results hold at the next-to-next-to-eikonal level with a proper definition of the leading eikonal piece, and can presumably be extended to leading logarithmic contributions at all orders in $(1-x)$. Analogous results are obtained for fragmentation functions in semi-inclusive e^+e^- annihilation.

One may ask to what extent the success of the present approach may be attributed, as suggested in [24, 32] for the splitting functions case, to the *classical nature* [36] of soft radiation. In fact, the main result of this paper for the (modified) DIS physical evolution kernel can be summarized (barring the δ -function contribution) by the following equation:

$$K(x, a_s, \lambda = 1) \sim \left[\frac{x}{1-x} + \frac{1}{2}(1-x) \right] \mathcal{J}((1-x)Q^2) + \text{subleading logarithms} , \quad (6.1)$$

where the second term (the “subleading logarithms”) is contributed by all powers in $(1-x)$ except the leading eikonal one. The first term in eq.(6.1) accounts for the leading logarithmic contributions to the modified kernel (together with some subleading logarithms) to *all* powers in $(1-x)$ at any given loop order, and implies leading logarithmic contributions are actually absent beyond $\mathcal{O}(1-x)$ power. This term has the remarkable effective one-loop splitting function form $4C_F a_{phys}((1-x)Q^2) p_{qq}(x)$, with the “physical coupling” $a_{phys}(Q^2) \equiv \frac{1}{4C_F} \mathcal{J}(Q^2)$.

As pointed out in [32], the $\frac{x}{1-x}$ part of the one-loop prefactor (eq.(2.8)) should be interpreted as corresponding to universal classical radiation, a QCD manifestation of the Low-Burnett-Kroll theorem [36], while the $1-x$ part represents a genuine quantum contribution. Now, it is clear that at the next-to-eikonal level, the $1-x$ part of the prefactor is irrelevant: only the “classical” $1/r$ part is required to separate those leading logarithms in the standard ($\lambda = 0$) physical evolution kernel which are correctly predicted in the present approach (the h_{ji} in eq.(4.1)), hence “inherited” in the sense of [32], from the “primordial” ones (those which at each loop order carry the *same* color factors as the leading $\mathcal{O}(1/(1-x))$ eikonal logarithms, and can thus be absorbed into the definition of the leading term). However, it appears from the results of section 4 that, at next-to-next-to-eikonal level, the full one-loop prefactor has to be used into the definition of the leading term

⁵Evolution equations involving similar kinematical rescaling factors have been suggested in the past: see e.g. eq.(3.2) in [20].

to properly isolate the “inherited” next-to-next-to-eikonal logarithms (the f_{ji}^c in eq.(4.1)). Moreover, although the “inherited” f_{ji}^c are purely “classical” (like the h_{ji}), the “inherited” g_{ji} at the $\mathcal{O}((1-x)^2)$ level are a mixture of “quantum” and “classical”. Indeed, setting $f_{10}^q = \frac{1}{2}k_{10}$ and $f_{21}^q = \frac{1}{2}k_{21}$ (the “quantum parts” of the $\mathcal{O}(1-x)$ coefficients), one finds $g_{21} = g_{21}^q + g_{21}^c$, with $g_{21}^q = k_{10}f_{10}^q = \frac{1}{2}k_{10}^2$ and $g_{21}^c = -\frac{1}{6}k_{10}^2$; and $g_{32} = g_{32}^q + g_{32}^c$, with $g_{32}^q = \frac{1}{2}k_{21}f_{10}^q + k_{10}f_{21}^q = \frac{3}{4}k_{21}k_{10}$ and $g_{32}^c = -\frac{1}{4}k_{21}k_{10}$. In both cases $g_{ji}^c = -\frac{1}{3}g_{ji}^q$, which shows the “inherited” g_{ji} coefficients are actually dominantly “quantum”.

It can be further checked [30] that the very same first term in eq.(6.1) also accounts for the leading logarithmic contributions to the $\lambda = -1$ modified SIA physical evolution kernel to *all* powers in $(1-x)$, which implies that the *leading logarithmic parts* of the *modified* DIS and SIA physical evolution kernels satisfy Gribov-Lipatov relation [37], namely we have:

$$K(x, a_s, \lambda = 1)|_{\text{LL}} = K_T(x, a_s, \lambda = -1)|_{\text{LL}} = p_{qq}(x) \mathcal{J}((1-x)Q^2)|_{\text{LL}} , \quad (6.2)$$

where $\mathcal{J}((1-x)Q^2)|_{\text{LL}} = A(a_s((1-x)Q^2))|_{\text{LL}} = \frac{A_1 a_s(Q^2)}{1+a_s(Q^2)\beta_0 L_x}$ is the leading logarithmic contribution to eq.(1.7). Indeed, once transformed back to the standard ($\lambda = 0$) physical kernels, eq.(6.2) is consistent with eq.(3.26) in [7] at least to next-to-next-to-eikonal order, and is probably correct to all orders in $(1-x)$ (with identically vanishing contributions beyond $\mathcal{O}(1-x)$ order). On the other hand, contrary to the splitting functions case where it has been checked up to three loops [32, 38], a full Gribov-Lipatov relation $K(x, a_s, \lambda = 1) = K_T(x, a_s, \lambda = -1)$ does not seem to hold for subleading logarithms beyond the leading eikonal level.

The resummation of the subleading logarithmic contributions at next-to-eikonal order in eq.(6.1), not addressed here, remains an open issue: the present method does not work for them, except in the conformal limit, where one recovers the results of [24].

Acknowledgements I thank G. Marchesini for stimulating discussions and hospitality at the University of Milano-Bicocca where part of this paper has been written. I also wish to thank the referees for constructive suggestions.

References

- [1] M. Kramer, E. Laenen and M. Spira, Nucl. Phys. B **511** (1998) 523 [arXiv:hep-ph/9611272].
- [2] G. Grunberg, arXiv:0710.5693 [hep-ph].
- [3] E. Laenen, L. Magnea and G. Stavenga, Phys. Lett. B **669** (2008) 173 [arXiv:0807.4412 [hep-ph]].
- [4] E. Laenen, G. Stavenga and C. D. White, JHEP **0903** (2009) 054 [arXiv:0811.2067 [hep-ph]].
- [5] S. Moch and A. Vogt, JHEP **0904** (2009) 081 [arXiv:0902.2342 [hep-ph]].
- [6] G. Grunberg and V. Ravindran, JHEP **0910** (2009) 055 [arXiv:0902.2702 [hep-ph]].
- [7] S. Moch and A. Vogt, JHEP **0911** (2009) 099 [arXiv:0909.2124 [hep-ph]].

- [8] G. Grunberg, arXiv:0910.3894 [hep-ph].
- [9] G. Soar, A. Vogt, S. Moch and J. Vermaseren, arXiv:0912.0369 [hep-ph].
- [10] W. Furmanski and R. Petronzio, *Z. Phys. C* **11** (1982) 293.
- [11] G. Grunberg, *Phys. Rev.* **D29** (1984) 2315.
- [12] S. Catani, *Z. Phys.* **C75** (1997) 665 [hep-ph/9609263].
- [13] J. Blumlein, V. Ravindran and W. L. van Neerven, *Nucl. Phys. B* **586** (2000) 349 [arXiv:hep-ph/0004172].
- [14] W. L. van Neerven and A. Vogt, *Nucl. Phys. B* **603** (2001) 42 [arXiv:hep-ph/0103123].
- [15] E. Gardi and G. Grunberg, *Nucl. Phys. B* **794** (2008) 61 [arXiv:0709.2877 [hep-ph]].
- [16] G. Sterman, *Nucl. Phys.* **B281** (1987) 310.
- [17] S. Catani and L. Trentadue, *Nucl. Phys.* **B327** (1989) 323.
- [18] G. P. Korchemsky and G. Marchesini, *Phys. Lett. B* **313** (1993) 433.
- [19] S. Catani, B. R. Webber and G. Marchesini, *Nucl. Phys. B* **349** (1991) 635.
- [20] D. Amati, A. Bassetto, M. Ciafaloni, G. Marchesini and G. Veneziano, *Nucl. Phys. B* **173** (1980) 429.
- [21] M. Ciafaloni, *Phys. Lett. B* **95** (1980) 113;
- [22] M. Ciafaloni and G. Curci, *Phys. Lett. B* **102** (1981) 352.
- [23] Y. L. Dokshitzer, V. A. Khoze and S. I. Troian, *Phys. Rev. D* **53** (1996) 89 [arXiv:hep-ph/9506425].
- [24] Yu. L. Dokshitzer, G. Marchesini and G. P. Salam, *Phys. Lett. B* **634** (2006) 504 [arXiv:hep-ph/0511302].
- [25] G. Altarelli and G. Parisi, *Nucl. Phys. B* **126** (1977) 298.
- [26] G. Curci, W. Furmanski and R. Petronzio, *Nucl. Phys. B* **175** (1980) 27.
- [27] S. Moch, J. A. M. Vermaseren and A. Vogt, *Nucl. Phys.* **B688** (2004) 101 [hep-ph/0403192].
- [28] B. Basso and G. P. Korchemsky, *Nucl. Phys. B* **775** (2007) 1 [arXiv:hep-th/0612247].
- [29] G. P. Korchemsky, *Mod. Phys. Lett. A* **4** (1989) 1257.
- [30] G. Grunberg, “On the large- x structure of physical evolution kernels”, to be published.
- [31] A. H. Mueller, *Nucl. Phys. B* **228** (1983) 351.
- [32] Yu. L. Dokshitzer and G. Marchesini, *Phys. Lett. B* **646** (2007) 189 [arXiv:hep-th/0612248].
- [33] M. Cacciari and S. Catani, *Nucl. Phys. B* **617** (2001) 253 [arXiv:hep-ph/0107138].
- [34] S. Moch and A. Vogt, *Phys. Lett. B* **680** (2009) 239 [arXiv:0908.2746 [hep-ph]].
- [35] S. Moch, J. A. M. Vermaseren and A. Vogt, *Nucl. Phys. B* **813** (2009) 220 [arXiv:0812.4168 [hep-ph]].
- [36] F. E. Low, *Phys. Rev.* **110** (1958) 974; T. H. Burnett and N. M. Kroll, *Phys. Rev. Lett.* **20** (1968) 86.
- [37] V. N. Gribov and L. N. Lipatov, *Sov. J. Nucl. Phys.* **15** (1972) 438, *ibid.* **15** (1972) 675.
- [38] A. Mitov, S. Moch and A. Vogt, *Phys. Lett. B* **638** (2006) 61 [arXiv:hep-ph/0604053].